

# Activity of Posaconazole versus Voriconazole for the Treatment of Invasive Aspergillosis in Adults Enrolled in a Clinical Trial

Mariana Castanheira<sup>1</sup>, Leah Woosley<sup>1</sup>, Mary Motyl<sup>2</sup>, Seongah Han<sup>2</sup>, Havilland Campbell<sup>2</sup>, Anjana Grandhi<sup>2</sup>, Hetty Waskin<sup>2</sup>  
<sup>1</sup>JMI Laboratories, North Liberty, IA, USA; <sup>2</sup>Merck & Co., Inc., Kenilworth, NJ, USA

## Introduction

- Invasive aspergillosis (IA) is a life-threatening disease with limited treatment options. IA is associated with delays in effective treatment and significant early mortality.
- Posaconazole is a broad-spectrum triazole antifungal that exhibits potent antifungal activity against a variety of yeasts and moulds.
- Posaconazole is approved by US-FDA for prophylaxis of invasive *Aspergillus* and *Candida* infections and for treatment of oropharyngeal candidiasis, including those infections refractory to itraconazole and/or fluconazole.
- We evaluated the antifungal susceptibility profiles of isolates collected during a randomized, prospective, phase 3, double-blind, double-dummy study comparing posaconazole with voriconazole given for ≤12 weeks in the primary treatment of IA (ClinicalTrials.gov, NCT01782131; EudraCT, 2011-003938-14) using CLSI and EUCAST reference testing methodologies.

## Materials and Methods

- A total of 127 isolates were recovered from documented infections during 2013 through 2019 from more than 90 medical centers located in 23 countries.
- Isolates were identified by DNA sequencing of 28S (all isolates) and 1 of the following genes: β-tubulin for *Aspergillus* spp., translation elongation factor (*TEF*) for *Fusarium* spp., or *ITS* for all other species of filamentous fungi and yeasts according to the CLSI MM18-A document.
- Susceptibility tests were conducted for posaconazole, itraconazole, voriconazole, caspofungin, and amphotericin B according to CLSI documents M27 (2017) and M38 (2018), as well as EUCAST document EDef 9.1.
- EUCAST clinical breakpoints (2018), CLSI breakpoints (M60, 2017), and epidemiological cutoff values (ECV; M59, 2018) were applied as interpretative criteria.
- Quality control (QC) was performed as recommended by CLSI and EUCAST guidelines using the following strains: *Candida parapsilosis* ATCC 22019, *Candida krusei* ATCC 6258, *Aspergillus flavus* ATCC 204304, *Aspergillus fumigatus* ATCC 204305, and *Aspergillus fumigatus* ATCC MYA-3626.

## Results

- Of the 127 samples tested, 119 were identified as *Aspergillus* species (Figure 1).
- Aspergillus fumigatus* (N=76) was the most prevalent species, followed by *A. flavus* species complex (N=19), *A. section Nigri* (N=10), and *A. section Terrei* (N=7; Figure 1).
- Overall, posaconazole (MIC<sub>50/90</sub>, 0.5/1 mg/L) displayed similar activity to voriconazole (MIC<sub>50/90</sub>, 0.5/1 mg/L) and itraconazole (MIC<sub>50/90</sub>, 1/2 mg/L) against 119 *Aspergillus* species isolates by both CLSI and EUCAST methods (Table 1).
- Posaconazole (MIC<sub>50/90</sub>, 0.5/0.5 mg/L) and voriconazole (MIC<sub>50/90</sub>, 0.25/0.5 mg/L) inhibited all 76 *A. fumigatus* isolates at an MIC of 1 mg/L (Figure 2).
- Among 19 *A. flavus* species complex isolates recovered from this study, posaconazole (MIC<sub>50/90</sub>, 0.5/1 mg/L), voriconazole (MIC<sub>50/90</sub>, 1/1 mg/L), and itraconazole (MIC<sub>50/90</sub>, 0.5/1 mg/L) displayed equivalent activity (Table 1 and Figure 2).
- Posaconazole (MIC<sub>50/90</sub>, 1/1 mg/L) also exhibited activity against 10 *A. section Nigri* and 7 *A. section Terrei* isolates (MIC<sub>50</sub>, 0.5 mg/L). Posaconazole activity was similar to the activity displayed by voriconazole and itraconazole (Table 1 and Figure 2).
- Azole activity against other moulds, such as Mucorales group isolates and *Fusarium incarnatum-equiseti* species complex, was limited (Table 2).

Figure 1 Distribution of main species and organism groups of fungi isolates





Figure 2 Posaconazole, voriconazole, and itraconazole MIC distributions against the main *Aspergillus* species



## Conclusions

- Posaconazole displayed good activity against all *Aspergillus* species isolates.
- Posaconazole *in vitro* activity against *Aspergillus* species was similar to that activity observed by voriconazole and itraconazole.
- Azoles exhibited limited activity against *Fusarium* spp. and Mucorales group isolates.

Table 1 Antimicrobial activity of posaconazole and comparator agents tested against *Aspergillus* spp. isolates

Antimicrobial agent (No. of isolates)	CLSI <sup>a</sup>				EUCAST <sup>b</sup>				
	MIC/MEC <sub>50</sub>	MIC/MEC <sub>90</sub>	%WT	%NWT	MIC <sub>50</sub>	MIC <sub>90</sub>	%S	%I	%R
<b><i>Aspergillus</i> spp. (n=119)</b>									
Amphotericin B	2	2			0.5	2			
Caspofungin	0.03	0.06			0.06	0.12			
Itraconazole	1	2			0.5	1			
Posaconazole	0.5	1			0.12	0.25			
Voriconazole	0.5	1			0.5	2			
<b><i>A. fumigatus</i> (n=76)</b>									
Amphotericin B	2	2	100	0	0.5	1	98.7	1.3	0
Caspofungin	0.03	0.06	100	0	0.06	0.12			
Itraconazole	1	1	94.7	5.3	0.5	0.5	100	0	0
Posaconazole	0.5	0.5			0.12	0.12	93.4	6.6	0
Voriconazole	0.25	0.5	100	0	0.5	1	94.7	5.3	0
<b><i>A. section Flavi</i> (n=19)</b>									
Amphotericin B	2	2	100	0	1	1			
Caspofungin	0.015		100	0	0.03	0.12			
Itraconazole	0.5	1	100	0	0.5	0.5	100	0	0
Posaconazole	0.5	1	84.2	15.8	0.12	0.25			
Voriconazole	1	1	100	0	1	2			
<b><i>A. section Nigri</i> (n=10)</b>									
Amphotericin B	0.5	1	100	0	0.25	0.25	100	0	0
Caspofungin	0.015		100	0	0.03	0.25			
Itraconazole	2	4	100	0	1	1			
Posaconazole	1	1	100	0	0.25	0.25			
Voriconazole	1	1	100	0	1	1			
<b><i>A. section Terrei</i> (n=7)</b>									
Amphotericin B	2		100	0	2				
Caspofungin	0.03		100	0	0.06				
Itraconazole	0.5		100	0	0.25		100	0	0
Posaconazole	0.5		100	0	0.12		100	0	0
Voriconazole	0.5		100	0	0.5				

<sup>a</sup> ECV criteria published in CLSI M59 (2018)  
<sup>b</sup> Clinical breakpoint criteria published by EUCAST (2019)  
 MIC = minimal inhibitory concentration, MEC = minimal effective concentration, WT = wild-type, NWT = non wild-type, S = susceptible, I = intermediate, R = resistant

## Acknowledgements

Funding for this research was provided by Merck Sharp & Dohme Corp., a subsidiary of Merck & Co., Inc., Kenilworth, NJ, USA. Merck was involved in the design and decision to present these results. Merck did not contribute to decisions in the collection, analysis, or interpretation of the data.

Table 2 Antimicrobial activity of posaconazole and comparator agents tested against 5 other mould isolates

Organism <sup>a</sup>	CLSI MIC/MEC (mg/L)				EUCAST MIC (mg/L)			
	Amp B	Itra	Posa	Vori	Amp B	Itra	Posa	Vori
<i>Rhizomucor pusillus</i>	1	0.5	0.5	0.25	0.25	0.25	0.06	0.5
<i>Rhizopus oryzae</i>	1	2	1	8	0.25	1	0.5	8
<i>Mucor circinelloides</i>	0.25	4	2	>8	0.06	>8	1	>8
<i>Lasiodiplodia</i> spp.	0.5	>8	8	2	>8	>8	>8	2
<i>Fusarium incarnatum-equiseti</i> species complex	2	8	2	2	0.5	>8	2	2

<sup>a</sup> Clinical breakpoint criteria or ECV interpretation unavailable  
 Amp B = amphotericin B, Itra = itraconazole, Posa = posaconazole, Vori = voriconazole

## References

Araujo, R., Oliveira, M., Amorim, A., & Sampaio-Maia, B. (2015). Unpredictable susceptibility of emerging clinical moulds to tri-azoles: review of the literature and upcoming challenges for mould identification. *Eur J Clin Microbiol Infect Dis*, 34: 1289–1301.

Noxafil® (posaconazole) package insert. Available at [https://www.access.data.fda.gov/drugsatfda\\_docs/label/2015/022003s018s020,0205053s002s004,0205596s001s003tbl.pdf](https://www.access.data.fda.gov/drugsatfda_docs/label/2015/022003s018s020,0205053s002s004,0205596s001s003tbl.pdf)

Clinical and Laboratory Standards Institute (2017). *M61Ed1E. Performance standards for antifungal susceptibility testing of filamentous fungi, first edition*. Wayne, PA: CLSI.

Clinical and Laboratory Standards Institute (2018). *M38Ed3. Reference method for broth dilution antifungal susceptibility testing of filamentous fungi, third edition*. Wayne, PA: CLSI.

Clinical and Laboratory Standards Institute (2020). *M59Ed3. Epidemiological cutoff values for antifungal susceptibility testing, second edition*. Wayne, PA: CLSI.

Clinical and Laboratory Standards Institute (2018). *MM18A Interpretive Criteria for Identification of Bacteria and Fungi by DNA Target Sequencing, Approved Guideline second edition*. Wayne, PA: CLSI.

EUCAST E.DEF 9.3.1 Method for the determination of broth dilution minimum inhibitory concentrations of antifungal agents for conidia forming moulds. January 2017. Available at [https://www.eucast.org/astoffungi/methods/antifungal susceptibility testing/ast\\_of\\_moulds/](https://www.eucast.org/astoffungi/methods/antifungal susceptibility testing/ast_of_moulds/)

EUCAST Clinical breakpoints for fungi v 9.0 - February 2018. Available at <https://www.eucast.org/astoffungi/clinicalbreakpointsforantifungals/>

Pfller MA, Woosley LN, Messer SA, Jones RN, Castanheira M. Significance of molecular identification and antifungal susceptibility of clinically significant yeasts and moulds in a global antifungal surveillance programme. *Mycopathologia*. 2012;174(4):259–271.

Contact Information:  
 Mariana Castanheira, PhD  
 JMI Laboratories  
 345 Beaver Creek Centre, Suite A  
 North Liberty, IA 52317  
 Phone: (319) 665-3370  
 Fax: (319) 665-3371  
 Email: [mariana-castanheira@jmlabs.com](mailto:mariana-castanheira@jmlabs.com)



<https://bit.ly/2Q3SuV4>